

D. Reporting templates

Based on the need to now report data not only for FC citywide but also separately for each of the four zones (NW, NE, SE, SW; broken down using College and Drake) the city has been divided into for the purpose of WNV surveillance and mosquito control, CSU had to generate a new reporting template. This new reporting template is tailored to provide data relevant to the Level III and IV entomological triggers (see below) for control measures in the “City of Fort Collins Program Response Guidelines to Mosquito Borne Arboviral Activity (July 2008 edition)”.

Level III

- Vector index > 0.5 and increasing
- *Culex* mosquito populations increasing and at or above historical average for that time period
- Mosquito infection rates of > 3.0 per thousand (0.3%) and increasing

Level IV

- Vector index > 0.75.
- *Culex* mosquito population above historical average for that time period
- Sustained mosquito infection rates of > 5.0 per thousand (0.5%)

The new reporting format comprises a set of 6 tables to address the current week (1a, 2a, 3a) and to provide seasonal and historical context (1b, 2b, 3b) (see full table formats on following pages)

- Table 1a. Vector Index for current week
- Table 1b. Vector Index for All *Culex* by week from June-August
- Table 2a. Vector abundance for current week
- Table 2b. Vector abundance for All *Culex* by week from June-August
- Table 3a. WNV infection rate per 1,000 females for current week
- Table 3b. WNV infection rate per 1,000 females for All *Culex* by week from June-August

Table 1a. Vector Index for current week

Week: 31	Mean abundance of females per trap night ¹		Estimate for proportion of females infected with WNV ²		Vector Index		
	<i>Cx. pipiens</i>	<i>Cx. tarsalis</i>	<i>Cx. pipiens</i>	<i>Cx. tarsalis</i>	<i>Cx. pipiens</i> ³	<i>Cx. tarsalis</i> ⁴	All <i>Culex</i> ⁵
FC – Zone NW	4.00	32.44	0.0000	0.0035	0.000	0.113	0.113
FC – Zone NE	9.70	59.90	0.0014	0.0034	0.014	0.207	0.220
FC – Zone SE	1.33	74.53	0.0061	0.0009	0.008	0.067	0.075
FC – Zone SW	2.56	10.11	0.0000	0.0000	0.000	0.000	0.000
FC – Citywide	4.09	48.84	0.0018	0.0020	0.007	0.096	0.104
LV	2.4	81.6	0.0000	0.0009	0.000	0.076	0.076
BE	0.5	6.3	0.0000	0.0000	0.000	0.000	0.000

¹From Table 2a (CDC light trap catches only).

²Derived from the data presented in Table 3a for estimated infection rate per 1,000 females (CDC light trap and gravid trap catches combined).

³Vector Index for *Cx. pipiens* = (Mean abundance of *Cx. pipiens* females per trap night) x (Estimate for proportion of all *Cx. pipiens* females infected with WNV).

⁴Vector Index for *Cx. tarsalis* = (Mean abundance of *Cx. tarsalis* females per trap night) x (Estimate for proportion of all *Cx. tarsalis* females infected with WNV).

⁵Vector Index for All *Culex* = (Vector Index for *Cx. pipiens*) + (Vector Index for *Cx. tarsalis*).

Table 1b. Vector Index for All *Culex* by week from June-August

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV	BE
	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹		
24	0.00	0.00	0.00	0.00	0.00	0.01	0.00	0.00	0.00	0.00	0.00	0.00
25	0.00	0.00	0.00	0.02	0.00	0.00	0.00	0.00	0.00	0.00	0.00	0.25
26	0.00	0.00	0.00	0.01	0.00	0.02	0.00	0.01	0.00	0.01	0.00	0.00
27	0.00	0.00	0.00	0.04	0.00	0.02	0.00	0.00	0.00	0.02	0.00	0.23
28	0.00	0.00	0.004	0.04	0.20	0.11	0.00	0.01	0.08	0.05	0.00	0.85
29	0.008	0.11	0.01	0.15	0.13	0.32	0.00	0.03	0.05	0.17	0.23	0.47
30	0.11	0.12	0.22	0.17	0.07	0.33	0.00	0.08	0.10	0.19	0.08	0.00
31		0.20		0.21		0.22		0.09		0.21		
32		0.16		0.34		0.39		0.11		0.27		
33		0.19		0.41		0.27		0.07		0.25		
34		0.14		0.15		0.31		0.05		0.18		
35		0.05		0.14		0.21		0.13		0.14		
36		0.03		0.04		0.00		0.00		0.02		
37		0.05		0.03		0.01		0.00		0.02		

¹ The historical average for Week 24-35 uses 2006-2015 historical data (2003-2005 surveillance data were excluded due to changes in trap locations from 2006 onwards). Weeks 36 & 37 historical average are calculated from 2015-2016 data (2015 is the first year to have surveillance data collected into weeks 36 & 37).

Table 2a. Vector abundance for current week (CDC light trap catches only)

Week: 31	Total number females collected			Number CDC light trap nights	Mean abundance of females per CDC light trap night		
	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>		Cx. <i>pipiens</i> ¹	Cx. <i>tarsalis</i> ²	All <i>Culex</i> ³
FC – Zone NW	36	292	328	9	4.00	32.44	36.44
FC – Zone NE	97	599	696	10	9.70	59.90	69.60
FC – Zone SE	20	1118	1138	15	1.33	74.53	75.87
FC – Zone SW	23	91	114	9	2.56	10.11	12.67
FC – Citywide	176	2100	2276	43	4.09	48.84	52.93
LV	93	3099	3192	38	2.45	81.55	84.00
BE	2	25	27	4	0.50	6.25	6.75

¹Mean abundance of *Cx. pipiens* females per CDC light trap night = (Total number *Cx. pipiens* females collected) / (Number CDC light trap nights).

²Mean abundance of *Cx. tarsalis* females per CDC light trap night = (Total number *Cx. tarsalis* females collected) / (Number CDC light trap nights).

³Mean abundance of All *Culex* females per CDC light trap night = (Total number All *Culex* females collected) / (Number CDC light trap nights).

Table 2b. Vector abundance for All *Culex* by week from June-August

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV	BE
	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹		
24	2.56	2.15	2.91	7.98	4.60	10.06	0.78	1.08	2.98	5.98	3.70	8.25
25	1.44	7.39	4.00	21.99	13.67	24.71	2.67	3.42	6.56	15.77	10.89	79.00
26	10.78	13.07	9.90	38.64	43.93	35.94	6.00	6.21	21.14	25.25	12.03	29.00
27	11.33	26.80	32.90	63.19	31.60	44.38	1.89	9.78	21.44	39.89	25.92	59.50
28	13.89	46.92	34.30	85.53	240.87	71.44	6.33	13.05	96.23	57.37	23.00	117.75
29	21.67	54.64	97.00	84.61	109.07	69.60	16.00	21.60	68.49	60.60	66.89	101.75
30	36.44	51.96	69.60	122.24	75.87	82.02	12.67	18.65	52.93	72.33	84.00	6.75
31		66.09		98.83		59.89		17.80		65.30		
32		43.19		70.40		52.87		15.11		47.39		
33		29.54		58.41		36.89		9.23		36.26		
34		19.16		41.75		24.61		6.80		25.01		
35		8.73		18.97		12.71		3.72		11.60		
36		11.00		22.15		6.73		3.11		10.45		
37		12.06		10.60		4.80		1.56		6.99		

¹The historical average for Week 24-35 uses 2006-2015 historical data (2003-2005 surveillance data were excluded due to changes in trap locations from 2006 onwards). Weeks 36 & 37 historical average are calculated from 2015-2016 data (2015 is the first year to have surveillance data collected into weeks 36 & 37).

Table 3a. WNV infection rate per 1,000 females for current week (CDC light trap and gravid trap catches combined)

Week: 31	Total number individuals examined			Total number pools examined			Total number WNV-infected pools			Estimate for WNV infection rate per 1,000 females ¹		
	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>	Cx. <i>pipiens</i>	Cx. <i>tarsalis</i>	All <i>Culex</i>
FC – Zone NW	171	292	463	9	13	22	0	1	1	0.00	3.49	2.19
FC – Zone NE	701	599	1300	19	16	35	1	2	3	1.43	3.45	2.39
FC – Zone SE	172	1118	1290	11	30	41	1	1	2	6.11	0.90	1.59
FC – Zone SW	91	91	182	8	10	18	0	0	0	0.00	0.00	0
FC – Citywide	1135	2100	3235	47	69	116	2	4	6	1.81	1.97	1.93
LV	11	1077	1088	3	25	28	0	1	1	0.00	0.93	0.92
BE	2	25	27	1	4	5	0	0	0	0.00	0.00	0

¹Maximum likelihood estimate (MLE) for WNV infection rate per 1,000 females calculated using the CDC PooledInfRate 4.0 plug-in for Excel.

Table 3b. WNV infection rate per 1,000 females for All *Culex* by week from June-August

Week	FC – Zone NW		FC – Zone NE		FC – Zone SE		FC – Zone SW		FC – Citywide		LV	BE
	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹	Current year	Historical average ¹		
24	0.00	0.00	0.00	0.00	0.00	1.25	0.00	0.00	0.00	0.48	0.00	0.00
25	0.00	0.00	0.00	0.62	0.00	0.00	0.00	0.00	0.00	0.07	0.00	3.18
26	0.00	0.00	0.00	0.11	0.00	0.43	0.00	0.44	0.00	0.27	0.00	0.00
27	0.00	0.00	0.00	0.24	0.00	0.44	0.00	0.00	0.00	0.31	0.00	3.83
28	0.00	0.00	1.57	0.50	0.79	0.87	0.00	0.40	0.85	0.56	0.00	7.18
29	3.30	1.67	0.81	1.25	1.05	2.81	0.00	0.96	1.09	1.83	3.46	4.98
30	2.19	2.53	2.39	1.89	1.59	4.47	0.00	2.94	1.93	2.99	0.92	0.00
31		4.03		2.18		4.64		3.33		3.42		
32		4.21		6.09		8.15		10.17		6.26		
33		7.56		7.12		10.13		5.50		8.29		
34		7.51		4.25		11.55		10.43		7.94		
35		9.70		5.99		15.63		45.89		11.12		
36		10.96		2.43		0.00		0.00		1.68		
37		2.21		2.52		1.85		0.00		3.01		

¹The historical average for Week 24-35 uses 2006-2015 historical data (2003-2005 surveillance data were excluded due to changes in trap locations from 2006 onwards). Weeks 36 & 37 historical average are calculated from 2015-2016 data (2015 is the first year to have surveillance data collected into weeks 36 & 37).

